



Media centre

Electromagnetic fields and public health

Electromagnetic hypersensitivity

Fact sheet N°296 December 2005

As societies industrialize and the technological revolution continues, there has been an unprecedented increase in the number and diversity of electromagnetic field (EMF) sources. These sources include video display units (VDUs) associated with computers, mobile phones and their base stations. While these devices have made our life richer, safer and easier, they have been accompanied by concerns about possible health risks due to their EMF emissions.

For some time a number of individuals have reported a variety of health problems that they relate to exposure to EMF. While some individuals report mild symptoms and react by avoiding the fields as best they can, others are so severely affected that they cease work and change their entire lifestyle. This reputed sensitivity to EMF has been generally termed "electromagnetic hypersensitivity" or EHS.

This fact sheet describes what is known about the condition and provides information for helping people with such symptoms. Information provided is based on a WHO Workshop on Electrical Hypersensitivity (Prague, Czech Republic, 2004), an international conference on EMF and non-specific health symptoms (COST244bis, 1998), a European Commission report (Bergqvist and Vogel, 1997) and recent reviews of the literature.

What is EHS?

EHS is characterized by a variety of non-specific symptoms, which afflicted individuals attribute to exposure to EMF. The symptoms most commonly experienced include dermatological symptoms (redness, tingling, and burning sensations) as well as neurasthenic and vegetative symptoms (fatigue, tiredness, concentration difficulties, dizziness, nausea, heart palpitation, and digestive disturbances). The collection of symptoms is not part of any recognized syndrome.

EHS resembles multiple chemical sensitivities (MCS), another disorder associated with low-level environmental exposures to chemicals. Both EHS and MCS are characterized by a range of non-specific symptoms that lack apparent toxicological or physiological basis or independent verification. A more general term for sensitivity to environmental factors is Idiopathic Environmental Intolerance (IEI), which originated from a workshop convened by the International Program on Chemical Safety (IPCS) of the WHO in 1996 in Berlin. IEI is a descriptor without any implication of chemical etiology, immunological sensitivity or EMF susceptibility. IEI

For more information contact:

WHO Media centre
Telephone: +41 22 791 2222
E-mail: mediainquiries@who.int

incorporates a number of disorders sharing similar non-specific medically unexplained symptoms that adversely affect people. However since the term EHS is in common usage it will continue to be used here.

Prevalence

There is a very wide range of estimates of the prevalence of EHS in the general population. A survey of occupational medical centres estimated the prevalence of EHS to be a few individuals per million in the population. However, a survey of self-help groups yielded much higher estimates. Approximately 10% of reported cases of EHS were considered severe.

There is also considerable geographical variability in prevalence of EHS and in the reported symptoms. The reported incidence of EHS has been higher in Sweden, Germany, and Denmark, than in the United Kingdom, Austria, and France. VDU-related symptoms were more prevalent in Scandinavian countries, and they were more commonly related to skin disorders than elsewhere in Europe. Symptoms similar to those reported by EHS individuals are common in the general population.

Studies on EHS individuals

A number of studies have been conducted where EHS individuals were exposed to EMF similar to those that they attributed to the cause of their symptoms. The aim was to elicit symptoms under controlled laboratory conditions.

The majority of studies indicate that EHS individuals cannot detect EMF exposure any more accurately than non-EHS individuals. Well controlled and conducted double-blind studies have shown that symptoms were not correlated with EMF exposure.

It has been suggested that symptoms experienced by some EHS individuals might arise from environmental factors unrelated to EMF. Examples may include “flicker” from fluorescent lights, glare and other visual problems with VDUs, and poor ergonomic design of computer workstations. Other factors that may play a role include poor indoor air quality or stress in the workplace or living environment.

There are also some indications that these symptoms may be due to pre-existing psychiatric conditions as well as stress reactions as a result of worrying about EMF health effects, rather than the EMF exposure itself.

Conclusions

EHS is characterized by a variety of non-specific symptoms that differ from individual to individual. The symptoms are certainly real and can vary widely in their severity. Whatever its cause, EHS can be a disabling problem for the affected individual. EHS has no clear diagnostic criteria and there is no scientific basis to link EHS symptoms to EMF exposure. Further, EHS is not a medical diagnosis, nor is it clear that it represents a single medical problem.

Physicians: Treatment of affected individuals should focus on the health symptoms and the clinical picture, and not on the person's perceived need for reducing or eliminating EMF in the workplace or home. This requires:

- a medical evaluation to identify and treat any specific conditions that may be responsible for the symptoms,
- a psychological evaluation to identify alternative psychiatric/psychological conditions that may be responsible for the symptoms,
- an assessment of the workplace and home for factors that might contribute to the presented symptoms. These could include indoor air pollution, excessive noise, poor lighting (flickering light) or ergonomic factors. A reduction of stress and other improvements in the work situation might be appropriate.

For EHS individuals with long lasting symptoms and severe handicaps, therapy should be directed principally at reducing symptoms and functional handicaps. This should be done in close co-operation with a qualified medical specialist (to address the medical and psychological aspects of the symptoms) and a hygienist (to identify and, if necessary, control factors in the environment that are known to have adverse health effects of relevance to the patient).

Treatment should aim to establish an effective physician-patient relationship, help develop strategies for coping with the situation and encourage patients to return to work and lead a normal social life.

EHS individuals: Apart from treatment by professionals, self help groups can be a valuable resource for the EHS individual.

Governments: Governments should provide appropriately targeted and balanced information about potential health hazards of EMF to EHS individuals, health-care professionals and employers. The information should include a clear statement that no scientific basis currently exists for a connection between EHS and exposure to EMF.

Researchers: Some studies suggest that certain physiological responses of EHS individuals tend to be outside the normal range. In particular, hyper reactivity in the central nervous system and imbalance in the autonomic nervous system need to be followed up in clinical investigations and the results for the individuals taken as input for possible treatment.

What WHO is doing

WHO, through its International EMF Project, is identifying research needs and co-ordinating a world-wide program of EMF studies to allow a better understanding of any health risk associated with EMF exposure. Particular emphasis is placed on possible health consequences of low-level EMF. Information about the EMF Project and EMF effects is provided in a series of fact sheets in several languages www.who.int/emf/.

FURTHER READING

WHO workshop on electromagnetic hypersensitivity (2004), October 25-27, Prague, Czech Republic, www.who.int/peh-emf/meetings/hypersensitivity_prague2004/en/index.html

COST244bis (1998) Proceedings from Cost 244bis International Workshop on Electromagnetic Fields and Non-Specific Health Symptoms. Sept 19-20, 1998, Graz, Austria

Bergqvist U and Vogel E (1997) Possible health implications of subjective

symptoms and electromagnetic field. A report prepared by a European group of experts for the European Commission, DGV. Arbete och Hälsa, 1997:19. Swedish National Institute for Working Life, Stockholm, Sweden. ISBN 91-7045-438-8.

Rubin GJ, Das Munshi J, Wessely S. (2005) Electromagnetic hypersensitivity: a systematic review of provocation studies. *Psychosom Med.* 2005 Mar-Apr;67(2):224-32

Seitz H, Stinner D, Eikmann Th, Herr C, Roosli M. (2005) Electromagnetic hypersensitivity (EHS) and subjective health complaints associated with electromagnetic fields of mobile phone communication---a literature review published between 2000 and 2004. *Science of the Total Environment*, June 20 (Epub ahead of print).

Staudenmayer H. (1999) *Environmental Illness*, Lewis Publishers, Washington D.C. 1999, ISBN 1-56670-305-0.